

SITE SYNOPSIS

SITE NAME: LOUGH CARRA/MASK COMPLEX

SITE CODE: 001774

This site is dominated by two large lakes, Lough Mask and Lough Carra, and includes the smaller Cloon Lough. On the western side, the site is overlooked by the Partry Mountains, while to the east the landscape is largely low-lying agricultural land. The nearest large town is Ballinrobe which is about 4 km east of Lough Mask. The general geological character of the area is Carboniferous limestones, with some shales and sandstones on the western side of Lough Mask.

The underlying geology results in a great diversity of habitats, which support many scarce and rare plants and animals. Six habitats which are on Annex I of the EU Habitats Directive are listed for this site, including two which are priority habitats - limestone pavement and Cladium fen.

Lough Mask, at over 8,000 ha, is the sixth largest lake in the country and with a maximum depth of 58 m it is one of the deepest. It is an excellent example of a lowland oligotrophic lake. Aquatic and wetland plant species present which are characteristic of this habitat include several Pondweed (*Potamogeton*) species, Water Lobelia (*Lobelia dortmanna*) and Shoreweed (*Littorella uniflora*). The eastern part of the lake is shallow and is edged by a lowlying shoreline which is subject to winter flooding. An intricate mixture of plant communities has developed on the limestone, with bare pavement, scrub-dominated pavement, dry grassland and heath. A variety of wetland habitats are also present, along with significant amounts of deciduous woodland along the eastern and southern shores. The western shoreline is less diverse and lacks the limestone communities. However, the fast-flowing Owenbrin River has created at its mouth an interesting delta of coarse sandy sediment.

Lough Carra, which is hydrologically linked to Mask, is one of the best examples in Ireland of a hard water marl lake. It is a shallow (mostly less than 2 m) predominantly spring fed lake with only a few streams flowing into it. Its well known pellucid green colour is due to calcareous encrustations. It has well developed stonewort communities in the submerged zones, with *Chara curta*, *C. desmacantha*, *C. rudis* and *C. contraria* recorded. Lough Carra, like the eastern and southern shores of Mask, is fringed by a diverse complex of limestone and wetland habitats.

The limestone pavement within this site represents the northern limit of the limestones of Clare and Galway. The limestone is variable in character, from open bare pavement to areas covered with dense scrub. Associated with the pavement are areas of dry calcareous grassland and dry heath. Characteristic species present include Bloody Crane's-bill (*Geranium sanguineum*), Yellow-wort (*Blackstonia perfoliata*), Carline Thistle (*Carlina vulgaris*), Blue Fleabane (*Erigeron acer*), Wild Madder (*Rubia peregrina*), Rustyback (*Ceterach officinarum*) and Quaking-grass (*Briza media*). Several plant species, notably Spring Gentian (*Gentiana verna*) and Dense-flowered Orchid (*Neotinea maculata*), occur at the northern limit of their distribution. The area is also noted for its diversity of orchid species. Scrub vegetation is variable in character, with extensive areas dominated by Hazel (*Corylus avellana*) and Hawthorn (*Crataegus monogyna*), with Buckthorn (*Rhamnus catharticus*), Alder Buckthorn (*Frangula alnus*), Spindle (*Euonymus europaeus*) and Ash (*Fraxinus excelsior*). The dry heath is well developed in places and is characterised by Gorse (*Ulex europaeus*), Bell Heather (*Erica cinerea*), Heather (*Calluna vulgaris*) and St. Dabeoc's Heath

(*Dabeocia cantabrica*). The diminutive orchid Lesser Twayblade (*Listera cordata*) occurs within the heath communities.

A wide range of wetland habitats occur around Lough Carra and along parts of the eastern and southern shores of Lough Mask, including *Cladium* fen and alkaline fen, both listed as Annex I habitats on the EU Habitats Directive. The *Cladium* occurs as pure stands in places but also intermixed with Black Bog-rush (*Schoenus nigricans*), Common Club-rush (*Scirpus lacustris*), Common Reed (*Phragmites australis*) and a number of sedge species (*Carex* spp.). The alkaline fens are more extensive than the *Cladium* fens and here Black Bog-rush is a dominant species. A rich diversity of flowering plant occurs in the fen communities. In addition to the fen habitats, there are sparse but widespread reed swamps, wet grassland and some freshwater marsh communities around the lake shores.

Broad-leaved deciduous woodland occurs fairly frequently around much of the shores of the lakes and on some of the islands. This is often scrub-type woodland, which may be either dry and dominated by Hazel, Hawthorn and Ash, or wet and dominated by Birch (*Betula* spp.), Willow (*Salix* spp.) and Alder (*Alnus glutinosa*). In places the woodland is more developed and includes Sessile Oak (*Quercus petraea*), Holly (*Ilex aquifolium*) and Rowan (*Sorbus aucuparia*).

A high concentration of rare plants are found at this site. Five species protected under the Flora Protection Order (1987) occur: Irish St. John's-wort (*Hypericum canadense*), Chives (*Allium schoenoprasum*), Pillwort (*Pilularia globulifera*), Irish Lady's-tresses (*Spiranthes romanzoffiana*), and Small Cudweed (*Logfia minima*). Two other Red Data Book plants, Alder Buckthorn (*Frangula alnus*) and Bird's-nest Orchid (*Neottia nidus-avis*), also occur, along with two Red Data Book Stonewort species - *Chara curta* and *Chara rudis*.

A large loft in the stable block of Curramore House provides a summer breeding site of the Lesser Horseshoe Bat (*Rhinolophus hipposideros*), a species listed on Annex II of the EU Habitats Directive. The bats gain access to the loft through windows that extend from the ground floor to the loft area. The building is surrounded by mixed woods and is close to the shores of Lough Mask; both of these habitats provide ideal foraging habitat for the bats. In 1993 more than 100 bats were counted at this site, which makes it of international importance. A second internationally important summer roost of Lesser Horseshoe Bats occurs within the site at Ballykyne, near Clonbur. Over 150 bats have been counted at this site in recent years.

The site provide excellent habitat for Otter (*Lutra lutra*), also an Annex II species on the Habitats Directive, and the area has Pine Marten (*Martes martes*), a species listed in the Irish Red Data Book.

The site has important bird interests, both in winter and summer. It provides feeding areas for part of the Erriff/Derrycraff population of Greenland White-fronted Geese. This flock has declined somewhat in recent years but is still of national importance, with an average spring peak from 1989-94 of 124 birds. The following count figures are the averages from surveys in January 1995 and January 1996: Wigeon 167; Mallard 397; Shoveler 57; Pochard 91; Tufted Duck 757; Goldeneye 158; Lapwing 233; Curlew 118. Also, 68 Whooper Swan and 25 Gadwall were recorded in January 1996. The Shoveler, Tufted Duck and Goldeneye populations are of national importance. Both lakes are traditional sites for breeding gulls and terns. In 1995, 44 pairs of Common Tern nested at Lough Mask, while in 1992 a census of gulls at both lakes resulted in the following: Black-headed Gull 1,451 pairs, Common Gull 407 pairs and Lesser Black-backed Gull 361 pairs. The Common Gull colony represents 11.3% of the national total, and the Lesser Black-backed Gull colony is 6.9% of the total.

The deep waters of Lough Mask are home to a population of the glacial relict Arctic Char (*Salvelinus alpinus*), and a rare shrimp (*Niphargus* spp.) is also found in these waters. Lough Mask is a very important Brown Trout fishery. White-clawed Crayfish (*Austropotamobius pallipes*), a species listed on Annex II of the Habitats Directive, has been recorded from Lough Carra.

This site is of considerable conservation importance as it has good examples of six habitats listed on Annex I of the EU Habitats Directive: lowland oligotrophic lakes, marl lakes, limestone pavement, dry heath, *Cladium* fen and alkaline fen. Some of these habitats are amongst the best examples of their kind in the country. It also has two Annex II mammal species and an Annex II invertebrate. The site is of ornithological importance for both wintering and breeding birds, with three Annex I Bird Directive species occurring regularly. A relatively large number of other nationally rare or localised plant and animal species occur, including the glacial relict Arctic Char.